

Powers of Edge Ideals with Linear Resolutions

Nursel Erey*

Abstract

We show that if G is a gap-free and diamond-free graph, then $I(G)^s$ has a linear minimal free resolution for every $s \geq 2$.

1 Introduction

Let $S = k[x_1, \dots, x_n]$ be a polynomial ring and let G be a finite simple graph with edge ideal $I(G)$. There has been recent interest in studying the asymptotic regularity of powers of edge ideals for various families of graphs [2, 3, 8, 10, 11]. It is known [9] that when I is a graded ideal, regularity of powers I^s is a linear function for large s . In particular, for any graph G , there exist constants c and s_0 such that $\text{reg}(I(G)^s) = 2s + c$ for all $s \geq s_0$. A special case of interest is to combinatorially characterize when the powers of edge ideals have eventually linear minimal free resolutions. Since the s th power of the edge ideal is generated in degree $2s$, the ideal $I(G)^s$ has a linear minimal free resolution if and only if $c = 0$ in the given formula.

It was proved by Francisco, Hà and Van Tuyl that if some power of $I(G)$ has linear minimal free resolution, then the complement graph G^c has no induced 4-cycles, i.e., G is gap-free. In this direction, Nevo and Peeva [11] raised the following question.

Question 1.1. Is it true that a graph G is gap-free if and only if $I(G)^s$ has a linear minimal free resolution for every $s \gg 0$?

This question is known to have a positive answer when the regularity of the edge ideal is 2. In fact, Herzog, Hibi and Zheng [8] proved that if $I(G)$ has a linear minimal free resolution, then so does $I(G)^s$ for all $s \geq 1$. Banerjee [2] proved that if G is a gap-free and cricket-free graph, then $I(G)^s$ has linear minimal free resolution for every $s \geq 2$. In this paper, we provide a new family of gap-free graphs for which the second and higher powers of edge ideals have linear minimal free resolutions.

*Department of Mathematics, North Dakota State University, Fargo, ND *e-mail*: nurselerey@gmail.com

2 Preliminaries

2.1 Graph theory

Throughout this paper we consider finite simple graphs, i.e., graphs with no loops or multiple edges. For any graph G we write $V(G)$ and $E(G)$ respectively for the vertex and edge sets of the graph. We say two vertices u and v are **adjacent** and write $uv \in G$, if there is an edge between them. The set $N(u) = \{v : v \text{ is adjacent to } u\}$ is called the **neighbor set** of u . Any $v \in N(u)$ is called a **neighbor** of u . A vertex is called an **isolated vertex** if it has no neighbors. A **complete graph** (or **clique**) is a graph such that every pair of vertices are adjacent. A complete graph on n vertices is denoted by K_n .

A graph H is called a **subgraph** of G if the vertex and edge sets of H are contained respectively in those of G . A subgraph H of G is called an **induced subgraph** if $uv \in G$ implies $uv \in H$ for every $u, v \in V(H)$. A subset A of the vertices of G is called **independent** if $uv \notin G$ for every $u, v \in A$. A **dominating clique** of a graph G is a complete subgraph H of G such that every vertex of G either belongs to H or is adjacent to some vertex of H .

For any graph K , we say G is **K -free** if K is not an induced subgraph of G . The **clique number** of G , denoted $\omega(G)$, is the number of vertices in a maximum clique of G . For any vertex v of G , the graph $G - v$ denotes the induced subgraph of G which is obtained from G by **removing the vertex** v . The **complement** of a graph G , denoted G^c , is a graph on the same vertex set such that for every vertex u and v , $uv \in G^c$ if $uv \notin G$.

Let F and G be graphs with no common vertices and let $v \in V(G)$. We say G' is obtained from G by **substituting** F for v if the vertex v is replaced by F and all the vertices of $N(v)$ are adjacent to all the vertices of F , i.e., $V(G') = (V(G) \cup V(F)) \setminus \{v\}$ and $E(G') = E(G - v) \cup E(F) \cup \{ab : a \in N(v), b \in V(F)\}$. For a vertex $v \in V(G)$ and a positive integer k , we say G' is obtained from G by **multiplying** v by k when G' is formed by substituting an independent set A of k vertices for v .

A **cycle** graph with vertices $v_1, \dots, v_n$ and edges $v_1v_2, \dots, v_{n-1}v_n, v_nv_1$ is denoted by $C_n = (v_1v_2 \dots v_n)$. The complement of a cycle graph is called an **anticycle**. A **path** in a graph G is a sequence of vertices $u_1, \dots, u_n$ such that $u_iu_{i+1} \in G$ for all $1 \leq i \leq n-1$. A **path graph** on n vertices, denoted by P_n , is a graph with n distinct vertices $u_1, \dots, u_n$ and $n-1$ edges $u_1u_2, u_2u_3, \dots, u_{n-1}u_n$. A graph which is isomorphic to the graph with vertices a, b, c, d and edges ab, bc, ac, ad, cd is called a **diamond**. A graph which is isomorphic to the graph with vertices w_1, w_2, w_3, w_4, w_5 and edges $w_1w_3, w_2w_3, w_3w_4, w_3w_5, w_4w_5$ is called a **cricket**.

A graph G is called **chordal** if every induced cycle in G has at most three vertices. A **bipartite** graph is a graph that does not contain any odd-length cycles. Two disjoint edges uv and ab of G are said to form a **gap** if the induced subgraph of G on the vertices u, v, a, b has only two edges. A graph G is called **gap-free** if no two edges form a gap, or equivalently, G^c is C_4 -free.

For any $v \in V(G)$ and any induced subgraph H of G , the **distance** between v and H is the number of edges in a shortest path from v to a vertex of H . In particular, the distance between v and H is zero when $v \in H$.

2.2 Castelnuovo-Mumford regularity of monomial ideals

Let $S = k[x_1, \dots, x_n]$ be a polynomial ring over a field k . The **Castelnuovo-Mumford regularity** of a monomial ideal I is given by

$$\operatorname{reg}(I) = \max\{j - i : b_{i,j}(I) \neq 0\}$$

where $b_{i,j}(I)$ are the graded Betti numbers of I . If G is a graph on the vertices $x_1, \dots, x_n$, then the **edge ideal** of G is defined as

$$I(G) = (xy : xy \text{ is an edge of } G).$$

The method of polarization reduces the study of minimal free resolutions of monomial ideals to that of square-free monomial ideals. Therefore quadratic monomial ideals can be studied via edge ideals. We recall the following well-known results.

Theorem 2.1. [6, Corollary 1.6.3] *Let $I \subseteq S$ be a monomial ideal and let I^{Pol} be its polarization. Then $\operatorname{reg}(I) = \operatorname{reg}(I^{\operatorname{Pol}})$.*

Lemma 2.2. *If $I \subseteq S$ is a monomial ideal, then $\operatorname{reg}(I, x) \leq \operatorname{reg}(I)$ for any variable x .*

We will make use of the following theorem to bound the regularity of monomial ideals.

Theorem 2.3. [7, Lemma 2.10] *Let $I \subseteq S$ be a monomial ideal, and let m be a monomial of degree d . Then*

$$\operatorname{reg}(I) \leq \max\{\operatorname{reg}(I : m) + d, \operatorname{reg}(I, m)\}.$$

Moreover, if m is a variable x appearing in I , then $\operatorname{reg}(I)$ is equal to one of these terms.

The next result is a restatement of Theorem 2.3 in terms of edge ideals. Note that for any vertex x of G , the set $\operatorname{st} x$ stands for $N(x) \cup \{x\}$.

Theorem 2.4. [7, Lemma 3.1] *Let x be a vertex of G with neighbors $y_1, \dots, y_m$. Then*

$$\operatorname{reg}(I(G)) \leq \max\{\operatorname{reg}(I(G - \operatorname{st} x)) + 1, \operatorname{reg}(G - x)\}.$$

Moreover, $\operatorname{reg}(I(G))$ is equal to one of these terms.

Banerjee [2] introduced the following total order on the minimal monomial generators of powers of edge ideals.

Definition 2.5. [2, Discussion 4.1] *Let I be an edge ideal which is minimally generated by the monomials $L_1, \dots, L_k$. Consider the order $L_1 > L_2 > \dots > L_k$ and let $n \geq 2$. For any minimal monomial generator M of I^n , the expression $M = L_1^{a_1} L_2^{a_2} \dots L_k^{a_k}$ is called **maximal** if for all expressions $M = L_1^{b_1} \dots L_k^{b_k}$, we have $(a_1, \dots, a_k) >_{\operatorname{lex}} (b_1, \dots, b_k)$. Given two minimal monomial generators M, N of I^n with maximal expressions $M = L_1^{a_1} L_2^{a_2} \dots L_k^{a_k}$ and $N = L_1^{c_1} L_2^{c_2} \dots L_k^{c_k}$ we set $M > N$ if $(a_1, \dots, a_k) >_{\operatorname{lex}} (c_1, \dots, c_k)$. We let $L^{(n)}$ denote the totally ordered set of minimal monomial generators of I^n .*

The following theorem follows from [2, Theorem 4.12].

Theorem 2.6. [2] Let $I = (L_1, \dots, L_k)$ be an edge ideal and let $s \geq 1$. Let $L^{(s)} : L_1^{(s)} > L_2^{(s)} > \dots > L_r^{(s)}$ be the order on the minimal monomial generators of I^s induced by the order $L_1 > \dots > L_k$ as described in Definition 2.5. Then for every $1 \leq \ell \leq r - 1$,

$$((I^{s+1}, L_1^{(s)}, \dots, L_\ell^{(s)}) : L_{\ell+1}^{(s)}) = (I^{s+1} : L_{\ell+1}^{(s)}), \text{ some variables}.$$

As a consequence of the theorem above, the following result was obtained in [2], which gives a sufficient condition for powers of edge ideals to have linear resolutions.

Theorem 2.7. [2, Corollary 5.3] Let G be a graph with edge ideal $I = I(G)$ and let $s \geq 1$. If $\text{reg}(I) \leq 4$ and $\text{reg}(I^{s+1} : m) \leq 2$ for every minimal monomial generator m of I^s , then I^{s+1} has linear minimal free resolution.

The following classical result of Fröberg [5] gives a combinatorial characterization of edge ideals which have linear minimal free resolutions.

Theorem 2.8. [5, Theorem 1] The minimal free resolution of $I(G)$ is linear if and only if the complement graph G^c is chordal.

2.3 Even-connection and gap-free graphs

In [2, Definition 6.2] the concept of even-connectedness was defined in order to describe in combinatorial terms the minimal generators of certain colon ideals of powers of edge ideals.

Definition 2.9. Let G be a graph. Two vertices u and v in G are said to be **even-connected** with respect to an s -fold product $e_1 \dots e_s$ of edges of G if there is a sequence of vertices $p_0, p_1, \dots, p_{2\ell+1}$ for some $\ell \geq 1$ in G such that

- (i) $p_i p_{i+1} \in G$ for all $0 \leq i \leq 2\ell$,
- (ii) $p_0 = u$ and $p_{2\ell+1} = v$,
- (iii) for all $0 \leq j \leq \ell - 1$, $p_{2j+1} p_{2j+2} = e_i$ for some i , and
- (iv) for all k , $|\{j \mid \{p_{2j+1}, p_{2j+2}\} = e_k\}| \leq |\{t \mid e_t = e_k\}|$.

In such case, $p_0, p_1, \dots, p_{2\ell+1}$ is called an **even-connection** between u and v with respect to $e_1 \dots e_s$.

Theorem 2.10. [2, Theorems 6.1 and 6.7] Let G be a graph with edge ideal $I = I(G)$ and let $s \geq 1$. Let $M = e_1 \dots e_s$ be a minimal monomial generator of I^s where $e_1, \dots, e_s$ are some edges of G . Then $(I^{s+1} : M)$ is minimally generated by monomials of degree 2, and uv is a minimal generator of $(I^{s+1} : M)$ if and only if either $uv \in G$ or u and v are even-connected with respect to M .

Consequently, if G is a graph and M is a minimal generator of $I(G)^s$ for some $s \geq 1$, then $(I(G)^{s+1} : M)^{\text{pol}}$ is the edge ideal of some associated graph G' . When G is a gap-free graph, some combinatorial properties of G' are summarized in the next result.

Theorem 2.11. [2, Lemmas 6.14 and 6.15] *Let G be a gap-free graph with edge ideal $I = I(G)$ and let $e_1, \dots, e_s$ be some edges of G where $s \geq 1$. Then the graph G' which is associated to $(I^{s+1} : e_1 \dots e_s)^{\text{pol}}$ is gap-free as well. Also if $C_n = (u_1 \dots u_n)$ is a cycle on $n \geq 5$ vertices such that C_n^c is an induced subgraph of G' , then C_n^c is an induced subgraph of G as well.*

The next theorem is one of the main graph theoretic tools that will be used in the sequel.

Theorem 2.12. [4, Theorems 2 and 3] *Let G be a gap-free graph with no isolated vertices.*

- (i) *If $\omega(G) \geq 3$, then G has a dominating clique on $\omega(G)$ vertices.*
- (ii) *If $\omega(G) = 2$ and G is not bipartite, then G can be obtained from a five-cycle by vertex multiplication.*

3 Colon ideals of powers of edge ideals

In this section, we first prove some graph theoretic results regarding the structure of (gap, diamond)-free graphs. Then we analyze the colon ideals of the form $(I(G)^{s+1} : m)$ where m is a minimal monomial generator of $I(G)^s$.

Proposition 3.1. *Let H be a graph which is obtained from G by multiplying a vertex $v \in V(G)$.*

- (i) *G is gap-free if and only if H is gap-free.*
- (ii) *G and H have the same clique number.*
- (iii) *If G is diamond-free, then H is diamond-free if and only if v does not belong to any triangle of G .*

Proof. Suppose that v is replaced with the independent set $\{v_1, \dots, v_k\}$.

(i) If H is gap-free, then G is gap-free since G can be considered as an induced subgraph of H . Assume for a contradiction G is gap-free but there is a gap between $ab, cd \in H$. Observe that since the vertices $v_1, \dots, v_k$ have the same neighbors in H , at most one of them belongs to $\{a, b, c, d\}$. Then this gives a gap in G .

(ii) Since every clique of H contains at most one vertex of $\{v_1, \dots, v_k\}$, the result follows.

(iii) Suppose that G is diamond-free. If v belongs to a triangle of G , then H clearly has an induced diamond. Conversely, suppose that H has an induced diamond with edges ab, ad, bd, bc, cd . Since G is diamond-free, without loss of generality we may assume that either $v_1 = a$ or $v_1 = d$. If $v_1 = a$, then (vbd) is a triangle in G . If $v_1 = d$, then (avb) is a triangle in G . $\square$

Lemma 3.2. *Let G be a gap-free and diamond-free graph with $\omega(G) \geq 3$ and with no isolated vertices. Then there is a dominating clique $K_{\omega(G)}$ such that every vertex outside of $K_{\omega(G)}$ is adjacent to exactly one vertex of $K_{\omega(G)}$. Moreover the following statements hold.*

- (i) *If $\omega(G) \geq 4$, then $G - K_{\omega(G)}$ is independent and in particular, G^c is chordal.*
- (ii) *If $\omega(G) = 3$, then for every vertex x of K_3 the set $N(x) \setminus V(K_3)$ is independent and in particular, $G - \text{st } x$ is bipartite.*

Proof. Let $K_{\omega(G)}$ be a dominating clique on the vertices $u_1, \dots, u_{\omega(G)}$ as in Theorem 2.12.

Suppose v is a vertex of G which is not in $K_{\omega(G)}$. Assume for a contradiction $vu_i \in G$ and $vu_j \in G$ for some $i \neq j$. Let k be different from i and j . Since G is diamond-free, $u_k v \in G$. Since k was arbitrary, it follows that v is connected to all vertices of $K_{\omega(G)}$. Thus G contains a clique with $\omega(G) + 1$ vertices, a contradiction.

(i) Let v and w be distinct vertices outside of $K_{\omega(G)}$. Assume for a contradiction $vw \in G$. Suppose $vu_i \in G$ and $wu_j \in G$ for some i, j . Since $\omega(G) \geq 4$, there exist distinct vertices u_k and u_ℓ such that $\ell \neq i, j$ and $k \neq i, j$. We already know that none of $vu_k, vu_\ell, wu_k, wu_\ell$ is in G . Hence there is a gap between vw and $u_k u_\ell$, a contradiction.

(ii) Let x, y and z be the vertices of K_3 . Assume for a contradiction $ux, vx, uv \in G$ for some $u, v \notin V(K_3)$. Since $ux \in G$, u cannot be adjacent to y or z . Similarly v cannot be adjacent to y or z . Hence there is a gap between uv and yz which is a contradiction. $\square$

Remark 3.3. Observe that the complement of a C_5 is again a cycle on 5 vertices. Therefore a graph is C_5 -free if and only if it is C_5^c -free. We shall use this fact for the rest of the paper without reference.

Lemma 3.4. *Suppose G is a gap-free graph.*

- (i) *If G is bipartite, then G^c is chordal.*
- (ii) *Let G be connected $(C_5, \text{diamond})$ -free and $\omega(G) = 3$. Then either $G = C_6^c$ or G^c is chordal.*

Proof. (i) Assume for a contradiction G is bipartite but G^c has an induced cycle $C_n = (x_1 x_2 \dots x_n)$ where $n \geq 4$. Since G is gap-free, we must have $n \geq 5$. Let $A \cup B$ be a bipartition of G where $x_2 \in A$. Then $x_4, x_5 \in B$ since $x_2 x_4, x_2 x_5 \in G$. As $x_1 x_4 \in G$ we get $x_1 \in A$. Similarly, as $x_1 x_3 \in G$, we get $x_3 \in B$. But then since $x_3 x_5$ is an edge of G , the set B is not independent, which is a contradiction.

(ii) Assume for a contradiction $G \neq C_6^c$ and G^c is not chordal. Then G^c has an induced $C_n = (x_1 \dots x_n)$ for some $n \geq 6$.

Case 1: Let $n = 6$. Then since $G \neq C_6^c$ and G is connected, without loss of generality assume that there is a vertex $y \in V(G) \setminus V(C_6^c)$ such that yx_1 is an edge of G . Since there is no gap between yx_1 and $x_2 x_6$, either yx_6 or yx_2 must be an edge of G . Without loss of generality assume that $yx_2 \in G$. Since the induced subgraph on the vertices x_1, x_3, x_6, x_2, y is not an induced 5-cycle of G , we must have either $x_3 y \in G$ or $x_6 y \in G$. Without loss

of generality suppose that $x_3y \in G$. Consider the induced subgraph H on the vertices x_1, x_3, x_5, y . If $yx_5 \notin G$, then H is an induced diamond, contradiction. If $yx_5 \in G$, then H is a clique on 4 vertices, which is again a contradiction as $\omega(G) = 3$.

Case 2: Let $n \geq 7$. Then the induced subgraph on the vertices x_1, x_3, x_4, x_6 is a diamond, contradiction. $\square$

Theorem 3.5. *Suppose G is both gap-free and diamond-free. Then $\text{reg}(I(G)) \leq 3$.*

Proof. If G is triangle-free, then it is cricket-free and thus the result follows from [2, Theorem 3.4]. If $\omega(G) \geq 4$, then the result follows from Lemma 3.2 and Theorem 2.8. Therefore we may assume that $\omega(G) = 3$. We proceed by induction on the number of vertices of G . If G is a triangle, then $\text{reg}(I(G)) = 2$. Let K_3 be a dominating clique as in Lemma 3.2 and let x be a vertex of K_3 . Since $G - x$ is either triangle-free or have clique number 3 we get $\text{reg}(I(G - x)) \leq 3$ by induction. From Lemma 3.2 and Lemma 3.4 the graph $(G - \text{st } x)^c$ is chordal and $\text{reg}(I(G - \text{st } x)) = 2$ follows from Theorem 2.8. Thus $\text{reg}(I(G)) \leq 3$ using Theorem 2.4. $\square$

Lemma 3.6. *Let G be a gap-free graph and let $e_1 \dots e_s$ be an s -fold product of edges. Let $C_n = (u_1 \dots u_n)$ be a cycle where $n \geq 5$. Suppose C_n^c is an induced anticycle in the graph which is associated to $(I(G)^{s+1} : e_1 \dots e_s)^{\text{pol}}$. Then $e_i \cap \{u_1, \dots, u_n\} = \emptyset$ for every $1 \leq i \leq s$.*

Proof. Let G' be the gap-free graph associated to $(I(G)^{s+1} : e_1 \dots e_s)^{\text{pol}}$. First note that by Theorem 2.11, C_n^c is also an induced anticycle in G . Let $e_i = xy$ be fixed. Notice that by symmetry, it is enough to consider the following two cases.

Case 1: Suppose $\{x, y\} \cap \{u_1, \dots, u_n\} = \{x\}$. Without loss of generality assume that $x = u_1$. Since there is no gap between xy and u_2u_n , either $yu_2 \in G$ or $yu_n \in G$. Without loss of generality suppose $yu_2 \in G$. Then u_2yxu_3 is an even-connection. Thus $u_2u_3 \in G'$ which is a contradiction.

Case 2: Suppose $\{x, y\} \cap \{u_1, \dots, u_n\} = \{x, y\}$. Without loss of generality, assume that $x = u_1$. Then $y = u_i$ for some $i \neq 2, n$. Since $n \geq 5$, either $i + 1 < n$ or $i > 3$. If $i + 1 < n$, then u_nyxu_{n-1} is an even-connection and $u_nu_{n-1} \in G'$, a contradiction. If $i > 3$, then u_2yxu_3 is an even-connection and $u_2u_3 \in G'$, a contradiction. $\square$

Lemma 3.7. *Let G be a (diamond, gap)-free graph and let $s \geq 1$. For every minimal generator m of $I(G)^s$, the graph associated to $(I(G)^{s+1} : m)^{\text{pol}}$ does not contain induced C_n^c for all $n \geq 6$.*

Proof. Let G' be the gap-free graph which is associated to $(I(G)^{s+1} : m)^{\text{pol}}$ as in Theorem 2.11. Assume for a contradiction G' has induced C_n^c for some $n \geq 6$ where $C_n = (v_1 \dots v_n)$. From Theorem 2.11, the anticycle C_n^c is an induced subgraph of G . Let $m = e_1 \dots e_s$ for some edges $e_1, \dots, e_s$ of G and let $e_1 = ab$. From Lemma 3.6 it follows that neither a nor b belongs to C_n . Since there is no gap between v_1v_3 and ab , without loss of generality assume that $v_1a \in G$. Notice that $v_2b \notin G$ since otherwise v_1abv_2 is an even-connection and $v_1v_2 \in G'$, which is a contradiction. Similarly, $v_nb \notin G$ as otherwise v_1abv_n would be even-connection and that would require $v_1v_n \in G'$. Since there is no gap

between v_2v_n and ab , without loss of generality assume that $v_2a \in G$. Observe that $v_3b \notin G$ as otherwise v_2abv_3 would be even connection and that would require $v_2v_3 \in G'$. Because there is no gap between v_3v_n and ab , either $v_na \in G$ or $v_3a \in G$. Without loss of generality, suppose $v_3a \in G$. If $av_5 \in G$, then the induced subgraph of G on the vertices v_2, v_3, v_5, a is a diamond, which is a contradiction. Otherwise, the induced subgraph of G on the vertices v_1, v_3, v_5, a is a diamond, again a contradiction. $\square$

Lemma 3.8. *Let G be a gap-free and diamond-free graph with $\omega(G) = 3$. Let $K_3 = (abc)$ be a dominating clique of G .*

- (i) *For all $s \geq 1$ and for all $(s-1)$ -fold product $e_1 \dots e_{s-1}$ edges of G , every induced C_5 in the graph of $(I(G)^{s+1} : (ab)e_1 \dots e_{s-1})^{\text{pol}}$ contains at least 2 vertices of K_3 .*
- (ii) *For all $s \geq 1$ and for all $(s-1)$ -fold product $e_1 \dots e_{s-1}$ edges of G , $\text{reg}((I(G)^{s+1} : (ab)e_1 \dots e_{s-1})^{\text{pol}}) = 2$.*

Proof. (i) Let $C_5 = (u_1u_2u_3u_4u_5)$ be an induced cycle of G' where G' is the graph associated to $(I(G)^{s+1} : (ab)e_1 \dots e_{s-1})^{\text{pol}}$ as in Theorem 2.11. Then since G is a subgraph of G' and $V(C_5) \subseteq V(G)$, the cycle C_5 is also an induced subgraph of G . Keeping Lemma 3.2 in mind, assume for a contradiction $V(C_5) \cap V(K_3)$ has at most one vertex.

Case 1: Suppose that $V(C_5) \cap \{a, b, c\} = \emptyset$. Since no set of 3 vertices of C_5 is independent, each of $N(a) \setminus V(K_3)$, $N(b) \setminus V(K_3)$ and $N(c) \setminus V(K_3)$ contains at least one vertex of C_5 . Without loss of generality assume that $(N(a) \setminus V(K_3)) \cap V(C_5) = \{u_5\}$, $(N(b) \setminus V(K_3)) \cap V(C_5) = \{u_1, u_3\}$ and $(N(c) \setminus V(K_3)) \cap V(C_5) = \{u_2, u_4\}$. Then, there is no edge in G that connects u_2u_3 and au_5 , which is absurd because G is gap-free.

Case 2: Suppose that $|V(C_5) \cap \{a, b, c\}| = 1$. Without loss of generality assume that $a = u_1 \in V(C_5) \cap \{a, b, c\}$. Then $(N(a) \setminus V(K_3)) \cap V(C_5) = \{u_2, u_5\}$. Without loss of generality we may assume that u_4 is adjacent to b and u_3 is adjacent to c in G . But then u_2abu_4 is an even-connection and $u_2u_4 \in G'$. This is a contradiction because C_5 is an induced cycle of G' .

(ii) Let G' be the gap-free graph which is associated to $(I(G)^{s+1} : (ab)e_1 \dots e_{s-1})^{\text{pol}}$ as in Theorem 2.11. From Fröberg's Theorem and Lemma 3.7 it suffices to show that G' has no induced C_5 . Assume for a contradiction G' has induced cycle $C_5 = (u_1 \dots u_5)$. From Lemma 3.6 we have $\{a, b\} \cap \{u_1, \dots, u_5\} = \emptyset$. On the other hand, by part (i) we have $|\{a, b, c\} \cap \{u_1, \dots, u_5\}| \geq 2$ which is a contradiction. $\square$

4 Regularity of powers of (diamond, gap)-free graphs

The authors of [1] classified all imperfect P_5 -free and diamond-free graphs based on a special family of graphs. Our main result Theorem 4.9 will be based on this classification. To state Arbib and Mosca's theorem, we need some notation first. All the graphs in [1, Fig. 3] contain an induced $C_5 = (u_0u_1u_2u_3u_4)$. Let D_k denote the set of points at distance k from this C_5 .

Theorem 4.1. [1, Theorem 1.6] Any connected $(P_5, \text{diamond})$ -free graph that properly contains an induced C_5 can be obtained from C_5 or from a graph among those of [1, Fig. 3] by multiplying some $v \in D_1 \cup C_5$ and/or substituting a P_3 -free graph for some $v \in D_2$.

Since any gap-free graph is P_5 -free we obtain the following corollary.

Corollary 4.2. Any connected $(\text{gap}, \text{diamond})$ -free graph that properly contains an induced C_5 can be obtained from C_5 or from a graph among those of Fig. 1 by multiplying some vertex which does not belong to a triangle.

Proof. First note that multiplying a vertex of a triangle yields a diamond. Observe that G_{10} and G_0 are the only graphs in [1, Fig. 3] that contain vertices of distance 2 from the induced $C_5 = (u_0 u_1 u_2 u_3 u_4)$. Observe that in [1, Fig. 3] there is a gap between the edges $u_0 a_0$ and $b_0 b_2$ of G_4 . Therefore a $(\text{gap}, \text{diamond})$ -free graph cannot be obtained from G_4 by Proposition 3.1. Let H be a graph that contains at least one edge. We claim that substituting H for y in G_{10} or G_0 yields a gap. Indeed, if $y_1 y_2$ is an edge of H , then there is a gap between $y_1 y_2$ and $u_1 u_2$. $\square$

Remark 4.3. Let H be a graph and let u be a vertex of H . Let G be obtained from H by multiplying the vertex u by the independent set U . If C_n is an induced cycle of G where $n \geq 5$, then C_n contains at most one vertex from U . In particular, if H is C_n -free where $n \geq 5$, then so is G .

The proof of the next lemma is computer aided. We used Maple to list the induced 5-cycles of graphs in Fig. 1.

Lemma 4.4. Let G be a $(\text{gap}, \text{diamond})$ -free graph which is obtained from a graph among the graphs $G_1, G_2, \dots, G_9$ in Fig. 1 by vertex multiplication. Let C_5 be an induced subgraph of G and $e = \{a, b\}$ be an edge of G such that $e \cap V(C_5) = \emptyset$. Then at least one of the following statements holds.

- (i) $e \in E(K_3)$ for some dominating clique K_3 of G .
- (ii) There exists distinct $u, v \in V(C_5)$ such that $au \in G, bv \in G$ and $uv \notin C_5$.

Proof. Using Corollary 4.2 and keeping Remark 4.3 in mind, we consider cases.

Case 1: Let G be obtained from G_1 by replacing the vertices u_0, u_1, u_4 respectively with the independent sets U_0, U_1, U_4 . Since $(u_0 \dots u_4)$ is the only induced 5-cycle of G_1 , we may assume that $C_5 = (u_0^1 u_1^1 u_2 u_3 u_4^1)$ for some $u_i^1 \in U_i$. Since e does not intersect C_5 , by the symmetry of the graph we may assume $e = u_0^2 u_4^2$ or $e = u_0^2 a_0$ for some $u_i^2 \in U_i \setminus \{u_i^1\}$. In both cases, one vertex of e is adjacent to u_3 and the other one is adjacent to u_1^1 . Since $u_1^1 u_3 \notin C_5$, the statement (ii) holds.

Case 2: Suppose that G is obtained from G_2 by replacing the vertex u_1 with $\{u_1^1, \dots, u_1^k\}$. Since $(u_0 \dots u_4)$ is the only induced 5-cycle of G_2 , we may assume that $C_5 = (u_0 u_1^1 u_2 u_3 u_4)$. But then every edge of G intersects C_5 .

Case 3: Suppose that G is obtained from G_3 by multiplying u_0, u_2, u_3, u_4 . For each $i = 0, 2, 3, 4$ let u_i be replaced with $\{u_i^1, \dots, u_i^{k_i}\}$ in G . Suppose e does not belong to the dominating clique $(b_0 b_2 u_1)$. Since the induced 5-cycles of G_3 are $(u_0 \dots u_4)$, $(b_0 u_1 u_2 u_3 u_4)$ and $(b_2 u_1 u_0 u_4 u_3)$, the symmetry of the graph allows us to consider the following cases.

Case 3.1: Suppose $C_5 = (b_0 u_1 u_2^1 u_3^1 u_4^1)$. Then e can have forms $u_0^i u_4^j, u_3^i u_4^j, u_3^i u_2^j$ or $u_3^i b_2$. If $e = u_0^i u_4^j$, then (ii) holds because the endpoints of e are respectively adjacent to u_1 and u_3^1 and $u_1 u_3^1 \notin C_5$. Similarly, if $e = u_3^i u_4^j$, then the endpoints of e are respectively adjacent to u_2^1 and b_0 but $u_2^1 b_0 \notin C_5$. Lastly, if $e = u_3^i u_2^j$ or $e = u_3^i b_2$ then the endpoints of e are respectively adjacent to u_4^1 and u_1 but $u_1 u_4^1 \notin C_5$.

Case 3.2: Suppose $C_5 = (u_0^1 u_1 u_2^1 u_3^1 u_4^1)$. Notice that if $e = u_p^i u_q^j$ for some $u_p u_q \in C_5$, then (ii) clearly holds. By the symmetry of the graph, we may assume that $e = b_0 u_4^i$. Then $u_3^1 u_4^i \in G$, $b_0 u_1 \in G$ but $u_1 u_3^1 \notin G$ and (ii) is satisfied.

Case 4: Suppose that G is obtained from G_5 by multiplying the vertices u_1, u_4 . Observe that $K_3 = (b_1 b_4 u_0)$ is a dominating clique and every induced 5-cycle of G contains the vertices u_0, u_2, u_3 . It follows that every edge of G either belongs to K_3 or intersects C_5 .

Case 5: Suppose that G is obtained from G_6 by multiplying the vertices u_1, u_4, b_2 . Suppose that u_1 and b_2 are respectively replaced with the independent sets U and B . Note that $(b_1 b_4 u_0)$ and $(a_0 u_2 u_3)$ are dominating cliques of G . Suppose that e does not belong to these dominating triangles. Note that every induced 5-cycle of G contains both u_0 and u_3 . So, e contains neither u_0 nor u_3 . Then e can take the forms $uu_2, ub, b_1 u_2$ or $b_1 b$ for some $u \in U, b \in B$. In each case, one vertex of e is adjacent to u_0 and the other one is adjacent to u_3 . But since $u_0 u_3 \notin C_5$, the statement (ii) holds.

Case 6: Suppose that G is obtained from G_7 by multiplying the vertices u_1, b_3, b_4 . Suppose that b_3 and b_4 are respectively replaced with the independent sets B_3 and B_4 . Note that $(a_2 u_0 u_4)$ and $(a_0 u_2 u_3)$ are dominating cliques of G . Suppose that e does not belong to these dominating triangles. Note that every induced 5-cycle of G contains both u_0 and u_2 . So, e contains neither u_0 nor u_2 . Then e can take the forms $u_3 u_4, u_3 b_4', b_3' u_4$ or $b_3' b_4$ for some $b_3' \in B_3, b_4' \in B_4$. In each case, one vertex of e is adjacent to u_0 and the other one is adjacent to u_2 . But since $u_0 u_2 \notin C_5$, the statement (ii) holds.

Case 7: Suppose that G is obtained from G_8 by multiplying the vertices b_2, b_4, u_1 . Suppose that u_1 and b_2 are respectively replaced with the independent sets U and B . Note that $(a_2 u_0 u_4)$ and $(a_0 u_2 u_3)$ are dominating cliques of G . Suppose that e does not belong to these dominating triangles. Note that every induced 5-cycle of G contains both u_0 and u_3 . So, e contains neither u_0 nor u_3 . Then e can take the forms $ub, uu_2, a_2 u_2$ or $a_2 b$ for some $b \in B, u \in U$. In each case, one vertex of e is adjacent to u_0 and the other one is adjacent to u_3 . But since $u_0 u_3 \notin C_5$, the statement (ii) holds.

Case 8: Since every vertex of G_9 belongs to a triangle, G_9 does not generate any other graphs. Let e be an edge which does not belong to any of the dominating triangles $(a_0 u_2 u_3)$, $(u_0 a_2 u_4)$ and $(b_0 b_2 u_1)$.

Case 8.1: Suppose C_5 contains a_0, a_2 and b_0 . Then e is equal to one of $u_3 u_4, u_3 b_2, u_2 u_1$ or $u_0 u_1$. If e is equal to $u_3 u_4$ or $u_3 b_2$, then one vertex of e is adjacent to a_0 and the other one is adjacent to a_2 . But $a_0 a_2 \notin C_5$ so (ii) holds. Otherwise, one vertex of e is adjacent

to a_2 and the other one is adjacent to b_0 and $b_0a_2 \notin C_5$.

Case 8.2: Suppose C_5 contains b_2, u_0 and u_3 . Then e is equal to one of b_0u_4, b_0a_0, a_2u_2 or u_1u_2 . If e is equal to b_0u_4 or b_0a_0 , then one vertex of e is adjacent to b_2 and the other one is adjacent to u_0 . But $b_2u_0 \notin C_5$ so (ii) holds. Otherwise, one vertex of e is adjacent to u_0 and the other one is adjacent to u_3 and $u_0u_3 \notin C_5$.

Case 8.3: Suppose C_5 contains u_1, u_2 and u_4 . Then e is equal to one of b_0a_0, u_0a_0, b_2a_2 or b_2u_3 . If e is equal to b_0a_0 or u_0a_0 , then one vertex of e is adjacent to u_4 and the other one is adjacent to u_2 . But $u_2u_4 \notin C_5$ so (ii) holds. Otherwise, one vertex of e is adjacent to u_1 and the other one is adjacent to u_4 . But $u_1u_4 \notin C_5$, completing the proof. $\square$




Figure 1: The list of graphs from which every (gap, diamond)-free graph in Corollary 4.2 can be generated by vertex multiplication.

Theorem 4.5. *Let G be a (gap, diamond)-free graph which is obtained from a graph among the graphs $G_1, G_2, \dots, G_9$ in Fig.1 by vertex multiplication. Then $\text{reg}(I(G)^{s+1}) = 2s + 2$ for*

all $s \geq 1$.

Proof. Since each of the graphs $G_1, G_2, \dots, G_9$ contains an induced C_5 and a triangle, all of them have clique number 3 by Lemma 3.2(i). Let s be fixed and let $m = e_1 \dots e_s$ be a monomial generator of $I(G)^s$ for some edges $e_1 := ab, e_2, \dots, e_s$ of G . We claim that $\text{reg}(I(G)^{s+1} : m) \leq 2$ and then proof follows from Theorem 3.5 and Theorem 2.7. Note that if e_1 is an edge of a dominating K_3 of G , then the proof follows from Lemma 3.8(ii). So, let us assume that e_1 does not belong to a dominating K_3 . Let G' be the gap-free graph which is associated to $(I(G)^{s+1} : m)^{\text{pol}}$ as in Theorem 2.11. From Fröberg's theorem and Lemma 3.7, it suffices to show that G' has no induced cycle on 5 vertices. Assume for a contradiction G' has an induced C_5 . Then C_5 is an induced subgraph of G by Theorem 2.11. We know that $e_1 \cap V(C_5) = \emptyset$ from Lemma 3.6. Now, from Lemma 4.4 there exists distinct $u, v \in V(C_5)$ such that $au \in G, bv \in G$ and $uv \notin C_5$. Then $uabv$ is an even connection with respect to $e_1 \dots e_s$ and $uv \in G'$. But then since C_5 is an induced subgraph of G' we get $uv \in C_5$, which is a contradiction completing the proof. $\square$

The proof of the theorem above does not work for (gap,diamond)-free graphs which are obtained from G_0 or G_{10} in Fig. 1. Using Macaulay 2, one can see $\text{reg}(I(G_0)^2 : ya_2) = 3$ and $\text{reg}(I(G_{10})^2 : a_0y) = 3$.

Proposition 4.6. *If G is a (gap,diamond)-free graph with $I = I(G)$ which can be generated from G_{10} in Fig. 1, then $\text{reg}(I^{s+1}) = 2s + 2$ for all $s \geq 1$.*

Proof. First note that from Corollary 4.2 the graph G is obtained from G_{10} by multiplying some of the vertices y, u_0, u_1, u_4 . Let $s \geq 1$ be fixed and let vertices y, u_0, u_1, u_4 of G_{10} be respectively replaced by the independent sets $\{y_1, \dots, y_k\}, \{u_0^1, \dots, u_0^{m_0}\}, \{u_1^1, \dots, u_1^{m_1}\}$ and $\{u_4^1, \dots, u_4^{m_4}\}$ in G . We first show that if $e_1 \dots e_s$ is an s -fold product of edges of G with the property that there exists $1 \leq i \leq s$ such that for all $1 \leq j \leq k$, $y_j \nmid e_i$, then the ideal $(I^{s+1} : e_1 \dots e_s)$ has regularity 2. To this end, suppose that $y_j \nmid e_1$ for all j .

From Theorems 2.1, 2.8, 2.11 and Lemma 3.7, it suffices to show that the graph G' which is associated to $(I^{s+1} : e_1 \dots e_s)^{\text{pol}}$ is C_5 -free. Assume for a contradiction G' has induced C_5 . Then from Theorem 2.11, C_5 is also an induced subgraph of G . Recalling Remark 4.3, observe that C_5 must contain both u_2 and u_3 . Therefore, without loss of generality, we may assume that $C_5 = (u_0^1 u_1^1 u_2 u_3 u_4^1)$. From Lemma 3.6 and the symmetry of the graph, we may assume that $e_1 = u_0^2 u_4^2$ or $e_1 = u_0^2 a_0$. If $e_1 = u_0^2 u_4^2$, then $u_3 u_4^2 u_0^2 u_1^1$ is an even-connection between u_3 and u_1^1 with respect to $e_1 \dots e_s$ and $u_3 u_1^1 \in G'$, which is a contradiction because C_5 is an induced subgraph of G' . Similarly, if $e_1 = u_0^2 a_0$, then $u_1^1 u_0^2 a_0 u_3$ is an even-connection between u_1^1 and u_3 with respect to $e_1 \dots e_s$ and $u_1^1 u_3 \in G'$, a contradiction. This completes the proof of our claim.

Consider the order $e_1 > \dots > e_r > e_{r+1} > \dots > e_{r+k}$ on the edges of G where $e_{r+j} = a_0 y_j$ for all $j = 1, \dots, k$. Let $M_1 > \dots > M_z > M_{z+1} > \dots > M_{z+t}$ be the order on the minimal monomial generators of I^s induced by the order on the edges as in Definition 2.5. Let z be the largest index such that M_z has an expression $M_z = e_{i_1} \dots e_{i_s}$ such that $y_j \nmid e_{i_1}$ for all $1 \leq j \leq k$. Observe that $M_{z+1}, \dots, M_{z+t}$ all have unique s -fold

product expressions up to the permutation of the edges in the product. It follows from the previously proved claim that

$$\text{reg}(I^{s+1} : M_i) = 2 \text{ for all } i = 1, \dots, z. \quad (1)$$

Then repeated use of Lemma 2.2, Theorem 2.6 and Theorem 2.3 yields

$$\begin{aligned} \text{reg}(I^{s+1}) &\leq \max\{\text{reg}((I^{s+1} : M_1)) + 2s, \text{reg}(I^{s+1}, M_1)\} \\ &\leq \max\{2 + 2s, \text{reg}((I^{s+1}, M_1) : M_2) + 2s, \text{reg}((I^{s+1}, M_1, M_2))\} \\ &= \max\{2 + 2s, \text{reg}((I^{s+1} : M_2), \text{some variables}) + 2s, \text{reg}((I^{s+1}, M_1, M_2))\} \\ &= \max\{2 + 2s, \text{reg}((I^{s+1}, M_1, M_2))\} \\ &\quad \vdots \\ &\leq \max\{2 + 2s, \text{reg}(I^{s+1}, M_1, \dots, M_z)\}. \end{aligned}$$

In order to repeat this process, it remains to show that for all $1 \leq i \leq t$,

$$\text{reg}((I^{s+1}, M_1, \dots, M_z, \dots, M_{z+i-1}) : M_{z+i}) \leq 2. \quad (2)$$

To this end, let $1 \leq i \leq t$ be fixed. Then $M_{z+i} = (y_{j_1}a_0)^{\alpha_1}(y_{j_2}a_0)^{\alpha_2} \dots (y_{j_q}a_0)^{\alpha_q}$ for some $\alpha_1, \dots, \alpha_q > 0$ and $k \geq j_1 > j_2 > \dots > j_q \geq 1$. Observe that $(I^{s+1} : M_{z+i}) = I$ and using Theorem 2.6 we obtain

$$((I^{s+1}, M_1, \dots, M_z, \dots, M_{z+i-1}) : M_{z+i}) = (I, \text{some variables}).$$

Notice that every variable $x \in \{u_0^1, u_0^2, \dots, u_0^{m_0}, u_2, u_3\}$ belongs to the variable generators of the ideal above. Indeed, for every such x , we have

$$M_{z+i-p} = (xa_0)(y_{j_1}a_0)^{\alpha_1-1}(y_{j_2}a_0)^{\alpha_2} \dots (y_{j_q}a_0)^{\alpha_q}$$

for some $p > 0$. Therefore we obtain

$$((I^{s+1}, M_1, \dots, M_z, \dots, M_{z+i-1}) : M_{z+i}) = (I, \text{some variables}, u_0^1, u_0^2, \dots, u_0^{m_0}, u_2, u_3).$$

Let H be the graph with edge ideal $I(H) = (y_1a_0, \dots, y_ka_0)$. Then

$$(I, \text{some variables}, u_0^1, u_0^2, \dots, u_0^{m_0}, u_2, u_3) = (I(H), \text{some variables}).$$

Since H^c is chordal, from Fröberg's Theorem $\text{reg}(I(H)) = 2$ and then Eq. (2) follows from Lemma 2.2. Now we have

$$\text{reg}(I^{s+1}) \leq \max\{2 + 2s, \text{reg}(I^{s+1}, I^s)\} = \max\{2 + 2s, \text{reg}(I^s)\}.$$

Hence the result follows by induction and Theorem 3.5. $\square$

Since the monomial ideals are polarized by their generators, the following follows from the construction.

Remark 4.7. If $I \subseteq S$ is a monomial ideal and $J \subseteq S$ is a square-free monomial ideal, then $(I + J)^{\text{pol}} = I^{\text{pol}} + J$.

Proposition 4.8. If G is a (gap, diamond)-free graph with $I = I(G)$ which can be generated from G_0 in Fig. 1, then $\text{reg}(I^{s+1}) = 2s + 2$ for all $s \geq 1$.

Proof. First note that from Corollary 4.2 the graph G is obtained from G_0 by multiplying some of the vertices u_1, y . Let $s \geq 1$ be fixed and suppose that the vertices u_1 and y of G_0 are respectively replaced by the independent sets $U = \{u_1^1, \dots, u_1^m\}$ and $Y = \{y_1, \dots, y_k\}$ in G . We claim that if $e_1 \dots e_s$ is an s -fold product of edges of G with the property that there exists $1 \leq i \leq s$ such that $x \nmid e_i$ for all $x \in U \cup Y$, then $\text{reg}(I^{s+1} : e_1 \dots e_s) = 2$.

To this end, suppose that $x \nmid e_1$ for all $x \in U \cup Y$. From Theorems 2.1, 2.8, 2.11 and Lemma 3.7, it suffices to show that the graph G' which is associated to $(I^{s+1} : e_1 \dots e_s)^{\text{pol}}$ is C_5 -free. Assume for a contradiction G' has induced C_5 . Then C_5 is also an induced subgraph of G because of Theorem 2.11. Observe that $(ya_2u_4u_3a_0)$ and $(u_0u_1u_2u_3u_4)$ are the only induced 5-cycles of G_0 . Because of Remark 4.3 we may assume without loss of generality that $C_5 = (y_1a_2u_4u_3a_0)$ or $C_5 = (u_0u_1^1u_2u_3u_4)$. If $C_5 = (y_1a_2u_4u_3a_0)$, then by Lemma 3.6, e_1 does not intersect any of a_2, u_4, u_3, a_0 and therefore e_1 contains a vertex from U , which is a contradiction. Similarly, if $C_5 = (u_0u_1^1u_2u_3u_4)$, then e_1 does not intersect any of u_0, u_2, u_3, u_4 and therefore e_1 contains a vertex from Y , which is a contradiction. This completes the proof of our claim.

Let $e_1, \dots, e_r$ be the edges of G which do not contain any vertex from $U \cup Y$. Consider the order

$$\begin{aligned} e_1 &> e_2 > \dots > e_r > y_1a_0 > y_1a_2 > y_2a_0 > y_2a_2 > \dots > y_ka_0 > y_ka_2 \\ &> u_1^1u_0 > u_1^1u_2 > u_1^2u_0 > u_1^2u_2 > \dots > u_1^mu_0 > u_1^mu_2 \end{aligned}$$

on the edges of G . Consider the order $M_1 > \dots > M_z > M_{z+1} > \dots > M_{z+t}$ on the minimal monomial generators of I^s induced by the order on the edges as in Definition 2.5. Let z be the largest index such that M_z has an expression $M_z = e_{i_1} \dots e_{i_s}$ such that $x \nmid e_{i_1}$ for all $x \in Y \cup U$. It follows from the previously proved claim that

$$\text{reg}(I^{s+1} : M_i) = 2 \text{ for all } i = 1, \dots, z. \quad (3)$$

Then repeated use of Lemma 2.2, Theorem 2.6 and Theorem 2.3 yields

$$\begin{aligned} \text{reg}(I^{s+1}) &\leq \max\{\text{reg}((I^{s+1} : M_1)) + 2s, \text{reg}((I^{s+1}, M_1))\} \\ &\leq \max\{2 + 2s, \text{reg}((I^{s+1}, M_1) : M_2) + 2s, \text{reg}((I^{s+1}, M_1, M_2))\} \\ &= \max\{2 + 2s, \text{reg}(((I^{s+1} : M_2), \text{some variables})) + 2s, \text{reg}((I^{s+1}, M_1, M_2))\} \\ &= \max\{2 + 2s, \text{reg}((I^{s+1}, M_1, M_2))\} \\ &\vdots \\ &\leq \max\{2 + 2s, \text{reg}(I^{s+1}, M_1, \dots, M_z)\}. \end{aligned}$$

In order to repeat this process, it remains to show that for all $1 \leq i \leq t$,

$$\text{reg}(J_i) := \text{reg}((I^{s+1}, M_1, \dots, M_z, \dots, M_{z+i-1}) : M_{z+i}) \leq 2. \quad (4)$$

Let $1 \leq i \leq t$ be fixed. Then M_{z+i} has a maximal expression where the exponent of xw is non-zero for some $x \in U \cup Y$ and $xw \in G$. Let $N(w) \setminus (U \cup Y) = \{w_1, \dots, w_\tau\}$. Then for every $1 \leq \kappa \leq \tau$ we have $M_{z+i}w_\kappa/x > M_{z+i}$ since $ww_\kappa \in G$ and $w, w_\kappa \notin U \cup Y$. Therefore from Theorem 2.1, Theorem 2.6 and Remark 4.7 it follows that

$$\begin{aligned} \text{reg}(J_i) &= \text{reg}(((I^{s+1} : M_{z+i}), w_1, \dots, w_\tau, \text{some variables})) \\ &= \text{reg}(((I^{s+1} : M_{z+i})^{\text{pol}}, w_1, \dots, w_\tau, \text{some variables})) \end{aligned}$$

Let G' be the gap-free graph which is associated to $(I^{s+1} : M_{z+i})^{\text{pol}}$ as in Theorem 2.11. Using Lemma 2.2 it suffices to show that $\text{reg}((I(G'), w_1, \dots, w_\tau)) \leq 2$. If $(I(G'), w_1, \dots, w_\tau)$ is generated in degree 1, we have nothing to show. So, let us assume that $((I(G'), w_1, \dots, w_\tau) = I(H)$ where $H = G' - \{w_1, \dots, w_\tau\}$ has at least one edge. By Fröberg's theorem and Lemma 3.7, it remains to show that H is C_5 -free. Assume for a contradiction C_5 is an induced cycle of H . Then C_5 is an induced cycle of G' since H is an induced subgraph of G' . From Theorem 2.11 it follows that C_5 is an induced cycle of G . Observe that every induced 5-cycle of G_0 contains both u_3 and u_4 . Therefore C_5 contains both u_3 and u_4 by Remark 4.3. If $w = a_0$ or $w = u_2$, then $u_3 \notin V(H)$, and $u_3 \notin V(C_5)$, a contradiction. Similarly, if $w = a_2$ or $w = u_0$, then $u_4 \notin V(H)$, and $u_4 \notin V(C_5)$, a contradiction.

Now we have

$$\text{reg}(I^{s+1}) \leq \max\{2 + 2s, \text{reg}(I^{s+1}, I^s)\} = \max\{2 + 2s, \text{reg}(I^s)\}.$$

Hence the result follows by induction and Theorem 3.5. $\square$

Finally, we prove the main result of this paper.

Theorem 4.9. *If G is a (gap, diamond)-free graph, then $\text{reg}(I(G)^s) = 2s$ for all $s \geq 2$.*

Proof. We may assume G is a connected gap-free and diamond-free graph since removal of isolated vertices does not change the edge ideal. If $\omega(G) \geq 4$, then the proof follows from Theorem 2.8, Lemma 3.2 and [8, Theorem 1.2]. If $\omega(G) < 3$, then G is cricket-free and the result follows from [2, Theorem 6.17]. Therefore let us assume that $\omega(G) = 3$. Note that C_6^c is cricket-free since C_6^c has no vertex with 4 neighbors. If G is C_5 -free, then the result follows from combining Theorem 2.8, Lemma 3.4, [2, Theorem 1.2] and [8, Theorem 3.2] because C_6^c is cricket-free. Therefore let us assume that G contains an induced cycle on 5 vertices. If G is obtained from C_5 by multiplying vertices, then $\omega(G) < 3$ by Proposition 3.1. Otherwise, the result follows from combining Corollary 4.2, Theorem 4.5, Proposition 4.6 and Proposition 4.8. $\square$

References

- [1] C. Arbib, R. Mosca, *On $(P_5, \text{diamond})$ -free graphs*, Discrete Math. 250 (2002), no. 1-3, 1–22.
- [2] A. Banerjee, *The regularity of powers of edge ideals*, J. Algebraic Combin. 41 (2015), no. 2, 303–321.
- [3] S. Beyarslan, H. T. Hà, T. N. Trung, *Regularity of powers of forests and cycles*, J. Algebraic Combin. 42 (2015), no. 4, 1077–1095.
- [4] F.R.K. Chung, A. Gyárfás, Z. Tuza, W.T. Trotter, *The maximum number of edges in $2K_2$ -free graphs of bounded degree*, Discrete Math. 81 (1990), no. 2, 129–135.
- [5] R. Fröberg, *On Stanley-Reisner rings*, Topics in algebra, Part 2 (Warsaw, 1988), 57–70, Banach Center Publ., 26, Part 2, PWN, Warsaw, 1990.
- [6] J. Herzog and T. Hibi, *Monomial Ideals*, Springer-Verlag, London, 2011.
- [7] H. Dao, C. Huneke, J. Schweig, *Bounds on the regularity and projective dimension of ideals associated to graphs*, J. Algebraic Combin. 38 (2013), no. 1, 37–55.
- [8] J. Herzog, T. Hibi, X. Zheng, *Monomial ideals whose powers have a linear resolution*, Math. Scand. 95 (2004), no. 1, 23–32.
- [9] V. Kodiyalam, *Asymptotic behaviour of Castelnuovo-Mumford regularity*, Proc. Amer. Math. Soc. 128 (2000), no. 2, 407–411.
- [10] M. Moghimian, S. A. Fakhari, S. Yassemi, *Regularity of powers of edge ideal of whiskered cycles*, Comm. Algebra 45 (2017), no. 3, 1246–1259.
- [11] E. Nevo, I. Peeva, *C_4 -free edge ideals*, J. Algebr. Combin. 37, 243–248 (2013).